

Study on Screening Length of Quark-AntiQuark Pair in a Hot Plasma of Two Dimensional Sphere

A. Nata Atmaja

*Research Center for Physics
Indonesian Institute of Sciences (LIPI)
Kompleks PUSPITEK Serpong
Tangerang 15310, Indonesia
ardian@teori.fisika.lipi.go.id*

ABSTRACT: We study screening length of a quark-antiquark pair moving in a plasma, corresponds to four dimensional AdS-Schwarzschild Black Hole, in AdS/CFT correspondence. The geodesic solution of the ends of string at the boundary reduces the problem to the motion in equatorial plane in which the separation length L of quark-antiquark pair is parallel to the angular velocity ω . The string configuration for quark-antiquark pair is parameterized by momentum transfer P which is related to momentum transfer in the drag force configuration. We compare the result by computing the screening length in quark-antiquark reference frame where the gravity dual is four dimensional Kerr-AdS Black Hole and we find a relation between mass parameters M_{Sch} of Schwarzschild-AdS Black Hole and M_{Kerr} of Kerr-AdS Black Hole in large mass limit.

KEYWORDS: [Screening Length](#), [Schwarzschild-AdS](#), [Kerr-AdS](#).

Contents

1. Introduction	1
2. Screening Length in Plasma Reference Frame	3
3. Screening Length in Quark-Antiquark Reference Frame	6
3.1 “Boost-AdS” Black Hole	6
3.2 Kerr-AdS Black Hole in Boyer-Linquist coordinate	7
3.3 Kerr-AdS Black Hole in Asymptotically AdS coordinate	7
4. Discussion	10

1. Introduction

One of the important signatures of Quark Gluon Plasma produced by heavy ion collision’s experiment at RHIC and the new LHC is suppression of J/ψ meson production, $c\bar{c}$ pair. This phenomena is understood qualitatively as the temperature of the plasma reaches above Hagedorn temperature the potential interaction of $c\bar{c}$ pair would not able to hold them anymore therefore J/ψ meson will dissociate and be screened in Quark Gluon Plasma [1]. The screening potential of $c\bar{c}$ pair has a dependence on separation length between c and $\bar{c}$, L . The length L could have value up to some maximum length L_s called screening length where beyond this length the screening potential becomes flat.

In string theory, heavy quarks pair described by Wilson loop is defined as an open string where both ends attached on the probe brane. Evaluating the Wilson loop will tell us information about screening length dependent of quark-antiquark potential. The procedure to evaluate the Wilson loop goes by extremizing the action of corresponding gravity dual theory [2, 3, 4].

A rigorous calculation on the screening length of a moving quark-antiquark pair at finite temperature four dimensional $\mathcal{N} = 4$ Supersymmetric Yang-Mills theory from AdS/CFT correspondence was pioneered by Liu, Rajagopal, and Wiedemann [5]. It was then generalized to arbitrary dimension for conformal and non-conformal gauge theories by Cáceres, Natsuume, and Okamura [6]. The calculations there were done by going to reference frame of the moving quark-antiquark pair or explicitly by boosting the background metric to direction of quark-antiquark’s velocity. A different approach was done in the reference frame of the plasma by Chernicoff,

Garcia, and Guijosa [7] and furthermore they also calculated the energy from both approaches. One interesting result of these calculations is the present of boost factor, $(1 - v^2)$, in the screening length. In [6], it was shown that the scale of this factor depends on the dimension of black hole backgrounds. However, this scaling factor is valid in ultrarelativistic limit and disagrees with numerical fitting that was found in [7]¹. For the case considered in this article both scalings coincide.

In this article, we are interested to compute the screening length for the case where the gauge theory lives in a compact two dimensional sphere S^2 or the corresponding background metric is four dimensional Schwarzschild-AdS Black Hole in Global coordinate. In Poincare coordinate case we can compute the screening length for arbitrary angle θ_L between the separation length of quark-antiquark pair and its velocity direction or hot wind plasma direction. Unfortunately the situation is quite restricted for Global coordinate case. Main reason is because the geodesic of external quarks at the boundary must follow the great circle solutions. Since we want to keep the screening length to be fixed, both quark and antiquark must stay on the same great circle plane. In that way, the only possible angle is at $\theta_L = 0$ or the separation length of quark-antiquark pair to be parallel with its angular velocity or the hot wind plasma. Furthermore, using $SO(3)$ symmetry we can rotate arbitrary great circle planes to equatorial plane.

In the next section 2, we will compute the screening length in the reference frame of the plasma. We plot numerically the screening length for various fixed angular velocities as a function of momentum conjugate of angular coordinate ϕ in space direction, $\pi_\phi^\sigma \equiv P$. In section 3, we proceed the computation in the reference frame of the quark-antiquark pair. Unlike the Poincare case we have more than one background metrics; we classify these into “Boost-AdS” and Kerr-AdS Black Holes. *A priori* we can not compare the results from these different backgrounds since there is no clear relation between some parameters of the background metrics. Using the fact that angular velocity ω and rotation parameter a are bounded, taking the scalar curvature is given by the same parameter l , and assuming the screening length is invariant under different coordinate representations, we are able to find relation between mass parameter M_{Sch} in Schwarzschild AdS Black Hole and mass parameter M_{Kerr} in Kerr-AdS Black Hole which is given by $M_{Kerr} = M_{Sch}(1 - a^2 l^2)^{3/2}$. We will plot the screening length as a function of angular velocity for these backgrounds and show that this mass parameters relation is in accordance with physical situation compared to the naive relation, $M_{Sch} = M_{Kerr}$. In the last section 4, we discuss the results in more detail.

¹The screening length for five dimensional Schwarzschild-AdS Black Hole in Poincare coordinate goes like $L_s \propto (1 - v^2)^{1/4}$, as derived in [6] and in agreement with [5], while in [7] it goes like $L_s \propto (1 - v^2)^{1/3}$.

2. Screening Length in Plasma Reference Frame

In the plasma reference frame the background metric is static and is given by four dimensional Schwarzschild-AdS Black Hole [8]

$$ds^2 = -r^2 h(r) dt^2 + \frac{1}{r^2 h(r)} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (2.1)$$

$$h(r) = l^2 + \frac{1}{r^2} - \frac{2M_{Sch}}{r^3}, \quad (2.2)$$

$$T_H = \frac{1}{4\pi} \left(\frac{1}{r_H} + 3r_H l^2 \right), \quad (2.3)$$

where l is the radius of curvature of AdS space, M_{Sch} is proportional to the mass of the Schwarzschild-AdS Black Hole, T_H is the Hawking temperature, and r_H is the even horizon given by the most positive real root of $h(r)$, $h(r_H) = 0$. The Schwarzschild-AdS Black Hole has a minimum temperature and two branches of high temperature. We choose the branch for $r_H l > 1/\sqrt{3}$ or $M_{Sch} l > \frac{2}{3\sqrt{3}}$ which is favoured thermodynamically [9].

Classical solution of a string is obtained by solving the equation of motion from the Nambu-Goto action under the background (2.1),

$$S = -\frac{1}{2\pi\alpha'} \int d\sigma^2 \sqrt{-\det g_{\alpha\beta}}, \quad (2.4)$$

$$g_{\alpha\beta} \equiv G_{\mu\nu} \partial_\alpha X^\mu \partial_\beta X^\nu,$$

where σ^α is the worldsheet coordinate, with $\sigma^\alpha \equiv (\tau, \sigma)$, $X^\mu(\sigma^\alpha)$ is the space-time coordinate where the string worldsheet is embedded, and $G_{\mu\nu}$ is the space-time metric (2.1). The equation of motion derived from action (2.4) is simply written as

$$\nabla_\alpha P_\mu^\alpha = 0, \quad (2.5)$$

$$P_\mu^\alpha = \frac{\pi_\mu^\alpha}{\sqrt{-g}}, \quad \pi_\mu^\alpha = \frac{\delta S}{\delta \partial_\alpha X^\mu},$$

where π_μ^α is the canonical worldsheet momentum with $g = \det g_{\alpha\beta}$.

In deriving the equation of motion we will use the gauge $\tau = t$ while for σ will be determined later for convenient. As it was explained in the Introduction we will consider the solution on equatorial plane $\theta = \pi/2$. Taking the ansatz

$$\phi(\sigma^\alpha) \equiv \omega\tau + \phi(\sigma), \quad (2.6)$$

$$r(\sigma^\alpha) \equiv r(\sigma), \quad (2.7)$$

for a moving quark-antiquark with angular velocity ω , the equations of motion are given by:

for r ,

$$\frac{1}{\sqrt{-g}} \left(\frac{\omega^2(3M_{Sch} - r)}{h^2 r^4} r'^2 + (-M_{Sch} + r + 2l^2 r^3) \phi'^3 \right) - \frac{\partial}{\partial \sigma} \left[\frac{r^3 h - \omega^2}{r^3 h \sqrt{-g}} r' \right] = 0, \quad (2.8)$$

and for ϕ ,

$$\partial_\sigma \pi_\phi^\sigma \equiv \frac{\partial}{\partial \sigma} \left[\frac{r^4 h}{\sqrt{-g}} \phi' \right] = 0, \quad (2.9)$$

with π_ϕ^σ is constant and $' \equiv \frac{\partial}{\partial \sigma}$. For quark-antiquark system, the solution for r must satisfy conditions

$$r(-L/2) = r(L/2) = +\infty, \quad r'(\sigma_p) = 0, \quad (2.10)$$

where the length of string is L and $-L/2 < \sigma_p < L/2$ is the turning point where r takes the minimum value $r(\sigma_p) = r_p > r_H$. Since we still have one degree of freedom to be gauged, we can take $\phi = \sigma$ and one can easily check that equations (2.8) and (2.9) are equivalent; it is so because we only have one function r to solve².

We are going to use equation (2.9) to find the separation length, which is now precisely L , of quark-antiquark pair. The solution for r is given by

$$r' = \pm \frac{r^4 h}{P} \sqrt{\frac{r^4 h - P^2}{r^4 h - \omega^2 r^4}}, \quad (2.11)$$

where $P \equiv \pi_\phi^\sigma$ is a constant denoting amount of ϕ -component of momentum flow or transfer on the string. Our convention is that $P \geq 0$ and positive sign in (2.11) corresponds to the momentum transfer from boundary to r_p and negative sign corresponds to the momentum transfer from r_p to boundary. The quark-antiquark pair can be built of two configurations (a) and (b) in Figure 1. It is necessary to take these two configurations in order for string to join up at $r = r_p$. Note that since ϕ coordinate is periodic, $0 \leq \phi < 2\pi$, there are two possible configurations of string for quark-antiquark pair with the separation length $0 \leq L \leq \pi/2$ and $\pi/2 < L < 2\pi$. For our purpose, it is natural to take string configuration with the shortest separation length.

Using symmetry, $\phi \rightarrow -\phi$, it is enough just to find solution of 2.11 with positive sign which corresponds to configuration (b) in Figure 1; this sets $\phi_p \equiv \sigma_p = 0$. At σ_p , we must have $r_p^4 h(r_p) - P^2 = 0$ and $r_p^4 h(r_p) - \omega^2 r_p^4 > 0$ which implies that $h(r_p) > 0$ or $r_p > r_H$ for $P \neq 0$. There is a critical radius r_c defined as $r_c^4 h(r_c) = \omega^2 r_c^4$ and it's also $r_c > r_H$ for $\omega \neq 0$. If $r_p = r_c$ then the condition (2.10) could not be satisfied and the string configuration would just be a single quark at the boundary where one

²We can simply choose a gauge $\phi = \omega\tau + \sigma$ in the first place which gives us the equation (2.8) that has more complicated form than equation (2.9).




Figure 1: Two different configurations determined by the sign in P . The positive sign in (b) corresponds to drag force configuration while negative sign (a) corresponds to unphysical energy solution [10, 11].

end of string at the boundary and the other end will be at the horizon. Therefore for quark-antiquark pair we must have $r_p > r_c$ or $P^2 > \omega^2 r_c^4$. Notice that this r_c equals to r_{Sch} appears in the drag force computation [12, 13]. We also interpret the constant π_ϕ^σ in [12, 13] equals to P because both are given by the same formula and a constant should be invariant under any gauge choice. So for some fixed angular velocity ω , or r_c , the string configuration is characterized by amount of momentum transfer P to horizon with $P^2 \geq \omega^2 r_c^4$. At $P = \omega r_c^2$, the string tends to represent a single quark and if we increase the momentum transfer P the quark-antiquark pair is formed. Figure 2 shows schematic pictures of string configuration describing quark-antiquark pair and drag force of a single quark.




Figure 2: Picture on the left shows at critical momentum transfer $P^2 = \omega^2 r_c^4$ the string describe a single quark while picture on the right is if we increase P such that $P^2 > \omega^2 r_c^4$.

From equation (3.8), we obtain an integral formula for computing separation length as below

$$\int_0^{L/2} d\sigma = \frac{L}{2} = \int_{r_p}^{\infty} \frac{P}{r^4 h} \sqrt{\frac{r^4 h - \omega^2 r^4}{r^4 h - P^2}}. \quad (2.12)$$

The integral above can not be solved analytically. So we plots numerically the separation length for various value of ω/l in Figure 3.




Figure 3: Plots of separation length for various $\omega/l = 0.1, 0.2, 0.4, 0.6, 0.8$ and with fixed $M_{Sch}l = 10$ as functions of Pl , with the corresponding colors are black, blue, green, brown, red.

As one can see, we produce similar behaviour of screening length L_s as in Poincare case. Our plots do not start from $Pl = 0$ which is different from the plots produced in [5], for $\theta_L = \pi/2$, since P is bounded from below $P > \omega r_c^2$.

3. Screening Length in Quark-Antiquark Reference Frame

A different way to compute screening length is by going to the reference frame of the moving quark-antiquark pair. In this case the metric will be seen as it is rotating at angular velocity proportional to angular velocity of the quark-antiquark pair. A usual procedure, as in Poincare case, of how to find the rotating metric is by boosting the static metric (2.1), at $\theta = \pi/2$, along ϕ -direction.

3.1 “Boost-AdS” Black Hole

In order to get the rotating metric at equatorial, we boost the static metric (2.1) with the following transformation

$$t \rightarrow \gamma \left(t - \frac{\omega}{l^2} \phi \right), \quad \phi \rightarrow \gamma (\phi - \omega t), \quad (3.1)$$

with $\gamma = \left(1 - \frac{\omega^2}{l^2}\right)^{-1/2}$ is the boost factor, and the resulting metric is

$$ds^2 = -r^2 h(r) \gamma^2 \left(dt - \frac{\omega}{l^2} d\phi\right)^2 + \frac{1}{r^2 h(r)} dr^2 + r^2 \gamma^2 (d\phi - \omega dt)^2. \quad (3.2)$$

Taking the same gauge, $t = \tau$ and $\phi = \sigma$, the static solution for radial coordinates, $r \equiv r(\sigma)$, is given by

$$r' = \frac{r^4 h}{P \gamma} \sqrt{\frac{r^4 h - P^2}{r^4 h - \omega^2 r^4}}, \quad (3.3)$$

where P is again the ϕ -component of momentum transfer as in previous static black hole case. One can immediately see from (3.3) that separation length will be different with the static case by a boost factor γ and hence the screening length as well, $L_{Boost} = \gamma L_{Sch}$.

3.2 Kerr-AdS Black Hole in Boyer-Linquist coordinate

The “Boost-AdS” metric in (3.2) is only valid at equatorial plane. In general there exist a solution for the full geometry namely Kerr-AdS Black Holes. There are many coordinates representation for Kerr-AdS [14]. The simplest one is using Boyer-Linquist coordinate. We assume the equatorial plane in Kerr-AdS is similar to equatorial plane in Schwarzschild-AdS. The four dimensional Kerr-AdS Black Hole metric in Boyer-Linquist coordinates at equatorial is given by [15, 16]

$$ds^2 = -\frac{\Delta_r}{r^2} \left(dt - \frac{a}{\Xi} d\phi\right)^2 + \frac{r^2}{\Delta_r} dr^2 + \frac{1}{r^2} \left(adt - \frac{r^2 + a^2}{\Xi} d\phi\right)^2, \\ \Delta_r = (r^2 + a^2)(1 + l^2 r^2) - 2M_{Kerr} r, \quad \Xi = 1 - a^2 l^2. \quad (3.4)$$

Using the gauge, $t = \tau$ and $\phi = \sigma$, the radial solution is now given by

$$r' = \frac{\Delta_r}{P \Xi^2} \sqrt{\frac{\Delta_r - P^2 \Xi^2}{\Delta_r - a^2}}. \quad (3.5)$$

The momentum transfer P here, although a constant, has a different formula with the one in Schwarzschild-AdS or “Boost-AdS” case and this could give us a different separation length function of P compare to the previous one. However since we only want to compute the screening length, maximum of separation length which is an invariant scalar under coordinate transformations, it would be natural to expect the same result as in the “Boost-AdS” case.

3.3 Kerr-AdS Black Hole in Asymptotically AdS coordinate

Another representation of Kerr-AdS metric is using Asymptotically AdS (AAdS) coordinate. Unlike the previous two rotating black hole metrics, the Kerr-AdS metric

in AAdS coordinate is AdS asymptotically, $Y \rightarrow \infty$, and hence is preferred by AdS/CFT correspondence prescription [17, 18]. In general, to get the full geometry of AAdS coordinate is quite involved, yet it is still possible for particular case such as at equatorial. The four dimensional Kerr-AdS metric in AAdS coordinates at equatorial can be obtained by the following coordinate transformation from metric (3.4) [15, 16]:

$$\Xi Y^2 = r^2 + a^2, \quad \Phi = \phi - al^2 t. \quad (3.6)$$

The resulting metric is

$$ds^2 = -\frac{\Delta_Y}{\Xi^2(\Xi Y^2 - a^2)} (dt - ad\Phi)^2 + \frac{\Xi^2 Y^2}{\Delta_Y} dY^2 + \frac{1}{\Xi Y^2 - a^2} (a(1 + l^2 Y^2)dt - Y^2 d\Phi)^2, \\ \Delta_Y = Y^2 \Xi^2 (1 + l^2 Y^2) - 2M_{Kerr} \sqrt{\Xi Y^2 - a^2}, \quad \Xi = 1 - a^2 l^2. \quad (3.7)$$

Using similar gauge as before, the radial solution, $Y \equiv Y(\sigma)$, is given by

$$Y' = \frac{\Delta_Y}{P\Xi^2} \frac{\sqrt{\Xi Y^2 - a^2}}{Y} \sqrt{\frac{\Delta_Y - P^2 \Xi^2}{\Delta_Y - a^2 \Xi^2 (1 + l^2 Y^2)^2}}. \quad (3.8)$$

Now, we are going to compute numerically and compare the screening length for all metrics (3.2), (3.4), and (3.7) as functions of angular velocity ω or a . To compare the results we need to find relations between all the black hole parameters. One immediately notice that all these metrics have the same curvature radius of AdS space which is l . Angular velocity ω in (3.2) is bounded at boundary by $l^2 \geq \omega^2$ while a is also bounded by $a^2 l^2 \leq 1$ and so we are tempted to take $\omega^2 = a^2 l^4$. Remaining parameter that needs to be related is the mass parameter. There is no clear relation between mass parameter of Schwarzschild-AdS Black Hole, M_{Sch} , and mass parameter of Kerr-AdS Black Hole, M_{Kerr} . One could naively take $M_{Sch} = M_{Kerr}$ and try to see if this is in accordance with our physical picture. We would expect the screening length calculated in the reference frame of quark-antiquark pair should be different by a factor of

$$\gamma = \frac{1}{\sqrt{1 - \frac{\omega^2}{l^2}}} \equiv \frac{1}{\sqrt{1 - a^2 l^2}} \quad (3.9)$$

compared to the screening length calculated in the reference frame of hot plasma. Furthermore, all screening length in reference frame of quark-antiquark pair should be given by the same plot.

As we can see from Figure 4, the screening length of Boyer-Linquist and AAdS Black Holes are much closer to the screening length of Schwarzschild-AdS Black Hole rather than to the screening length of “Boost-AdS” Black Hole. Hence its is unlikely that M_{Sch} equals to M_{Kerr} and so we need to find how they are related exactly.




Figure 4: Plots of screening length, with $M_{Sch} = M_{Kerr} = 30/l$, for different black holes: Schwarzschild-AdS (Black), “Boost-AdS” (Red), Boyer-Linquist (Blue), and Asymptotically AdS (Green).

Suppose that Y_p , which is solution to $\Delta_Y(Y_p) = P^2\Xi^2$, is very large such that $l^2Yp^2 \gg 1$ then $1 + l^2Yp^2 \approx l^2Yp^2$. The equation (3.8) now becomes

$$Y' \approx \frac{l^2Y^4 - 2M_{Kerr}\Xi^{-3/2}Y}{P} \Xi^{1/2} \sqrt{\frac{l^2Y^4 - 2M_{Kerr}\Xi^{-3/2}Y - P^2}{l^2Y^4 - 2M_{Kerr}\Xi^{-3/2}Y - a^2l^4Y^4}}. \quad (3.10)$$

Taking large r_p in (3.3) such that $l^2r_p^2 \gg 1$, the equation is now given by

$$r' \approx \frac{l^2r^4 - 2M_{Sch}r}{\gamma P} \sqrt{\frac{l^2r^4 - 2M_{Sch}r - P^2}{l^2r^4 - 2M_{Sch}r - a^2l^4r^4}}, \quad (3.11)$$

with $\omega^2 = a^2l^4$. Comparing both equations (3.10) and (3.11), we can extract a relation

$$M_{Kerr} = M_{Sch}(1 - a^2l^2)^{3/2}. \quad (3.12)$$

The conditions for large r_p and Y_p can be satisfied if we take a limit of large M_{Sch} and M_{Kerr} . Recall that this is also the same limit if one would like to go from Global coordinate to Poincare coordinate of AdS [8]. Using this relation we obtain the numerical plots in Figure 5 where now the screening length of Kerr-AdS Black Holes closer to the screening length of “Boost-AdS” Black Hole which is in accordance with the physical picture we expected before.




Figure 5: Plots of screening length, with $M_{Sch} = \gamma^3 M_{Kerr} = 30/l$, for different black holes: Schwarzschild-AdS (Black), “Boost-AdS” (Red), Boyer-Linquist (Blue), and Asymptotically AdS (Green).

4. Discussion

In section 2, we have computed the separation length as a function of momentum transfer P and plotted it for various angular velocity ω . In the plots of Figure 3, there is minimum value of separation length for low momentum transfer P . This is because P is bounded from below where its value has to be bigger than the drag force momentum transfer, $P = \omega r_c^2$. Below this value there is no physical solution for string configuration of quark-antiquark pair. Therefore right after the momentum transfer P passes this bound the quark-antiquark pair is formed at the boundary with non-zero separation length but we suspect that this quark-antiquark pair is unstable. The reason is because the tip of the string in this configuration, located at r_p , is very close to r_c ; we identify r_c as a “shifted” horizon where at $\omega = 0$, $r_c = r_H$. However, there is another configuration of string, which is much more stable than the previous one, with the same separation length at high momentum transfer P where the tip of string, r_p , is far away from r_c . We then speculate that momentum transfer P in the string configuration of quark-antiquark pair is related to the strength of potential interaction between quark and anti-quark. It is interesting to study in more detail how exactly the relation is.

We also computed the screening length in the reference frame of quark-antiquark pair. Although there are more than one background metrics, we argued that the resulting screening length should not be different significantly. To compare the result for these backgrounds, we found a relation between parameters of this black hole backgrounds. At the end, we found that $M_{Kerr} = M_{Sch}(1 - a^2)^{3/2}$ is more suitable

for our physical picture at large mass limit.

In the literature, we have not found so far the relation between mass parameter of Schwarzschild-AdS Black Hole and mass parameter of Kerr-AdS Black Hole. A general procedure for constructing the metric of Kerr-AdS Black Holes in [14] was built of AdS metric plus the mass term, scaled with mass parameter of Kerr-AdS, in the Kerr-Schild form. Thus it has no direct connection with the mass parameter of Schwarzschild-AdS Black Hole. It would be interesting to find a procedure for constructing Kerr-AdS Black Hole from Schwarzschild-AdS Black Hole using the mass relation obtained in this article.

Acknowledgments

I am grateful to KEK group theory for hospitality and in particular to Makoto Natsuume for useful discussion during the initial work of this article.

References

- [1] T. Matsui and H. Satz, “J/psi Suppression by Quark-Gluon Plasma Formation,” *Phys. Lett. B* **178**, 416 (1986).
- [2] J. M. Maldacena, “Wilson loops in large N field theories,” *Phys. Rev. Lett.* **80**, 4859 (1998) [hep-th/9803002].
- [3] S. J. Rey and J. T. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity,” *Eur. Phys. J. C* **22**, 379 (2001) [arXiv:hep-th/9803001].
- [4] S. -J. Rey, S. Theisen and J. -T. Yee, “Wilson-Polyakov loop at finite temperature in large N gauge theory and anti-de Sitter supergravity,” *Nucl. Phys. B* **527**, 171 (1998) [hep-th/9803135].
- [5] H. Liu, K. Rajagopal and U. A. Wiedemann, “An AdS/CFT Calculation of Screening in a Hot Wind,” *Phys. Rev. Lett.* **98**, 182301 (2007) [hep-ph/0607062].
- [6] E. Caceres, M. Natsuume and T. Okamura, “Screening length in plasma winds,” *JHEP* **0610**, 011 (2006) [hep-th/0607233].
- [7] M. Chernicoff, J. A. Garcia and A. Guijosa, “The Energy of a Moving Quark-Antiquark Pair in an N=4 SYM Plasma,” *JHEP* **0609**, 068 (2006) [hep-th/0607089].
- [8] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” *Adv. Theor. Math. Phys.* **2**, 505 (1998) [arXiv:hep-th/9803131].
- [9] S. Hemming and L. Thorlacius, “Thermodynamics of Large AdS Black Holes,” *JHEP* **0711**, 086 (2007) [arXiv:0709.3738 [hep-th]].

- [10] S. S. Gubser, “Drag force in AdS/CFT,” *Phys. Rev. D* **74**, 126005 (2006) [arXiv:hep-th/0605182].
- [11] C. P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L. G. Yaffe, “Energy loss of a heavy quark moving through $N = 4$ supersymmetric Yang-Mills plasma,” *JHEP* **0607**, 013 (2006) [arXiv:hep-th/0605158].
- [12] A. Nata Atmaja and K. Schalm, “Anisotropic Drag Force from 4D Kerr-AdS Black Holes,” *JHEP* **1104**, 070 (2011) [arXiv:1012.3800 [hep-th]].
- [13] S. Chunlen, K. Peeters and M. Zamaklar, “Finite-size effects for jet quenching,” arXiv:1012.4677 [hep-th].
- [14] G. W. Gibbons, H. Lu, D. N. Page and C. N. Pope, “The general Kerr-de Sitter metrics in all dimensions,” *J. Geom. Phys.* **53**, 49 (2005) [arXiv:hep-th/0404008].
- [15] S. W. Hawking, C. J. Hunter and M. Taylor, “Rotation and the AdS/CFT correspondence,” *Phys. Rev. D* **59**, 064005 (1999) [arXiv:hep-th/9811056].
- [16] G. W. Gibbons, M. J. Perry and C. N. Pope, “The first law of thermodynamics for Kerr - anti-de Sitter black holes,” *Class. Quant. Grav.* **22**, 1503 (2005) [arXiv:hep-th/0408217].
- [17] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
- [18] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428**, 105 (1998) [arXiv:hep-th/9802109].